PROFESSIONALS

Maintaining client/patient safety is paramount when counseling someone who may be suicidal. Treat all suicide expressions with great importance. Screen for suicidal thoughts or plans with anyone who appears seriously depressed, makes suicidal references or has a history of attempts. Develop a safety and risk management process with client that would include the client's commitment to treatment, willingness to follow advice and removal of means. Do not rely solely on "No Suicide Contracts". Assess lethality of client's medication regimen. Assess client's risk of harm to others. Provide accessibility 24 hours a day until psychiatric referral can be realized.

Screening Tool- "IS PATH WARM" from American Association of Suicidology

- <u>I</u>DEATION-threatened or communicated
- <u>S</u>UBSTANCE ABUSE-excessive or increased
- PURPOSELESS-no reason for living
- ANXIETY-agitation or insomnia
- <u>TRAPPED-feeling there's no way out</u>
- HOPELESSNESS

- <u>W</u>ITHDRAWAL-from friends, family, society
- <u>A</u>NGER-uncontrolled, rage, seeking revenge
- <u>R</u>ECKLESSNESS-risky behaviors, actions, unthinking
- MOOD CHANGES-dramatic

Resources:

Nassau SPEAKS Out Resources

Suicide Assessment Five-Step Evaluation and Triage SAFE-T

American Association of Suicidology

Suicide Prevention Resource Center

Suicide Prevention among LGBT Youth

Evidence-based Curricula, i.e., ASIST, The Connect, Creating Suicide Safer Schools, SuicideTALK

CDC Suicide and Violence Prevention

Program to Encourage Active Rewarding Lives for Seniors (PEARLS)

Suicide Prevention Training for Gatekeepers of Older Adults

Role of Managers in Preventing Suicide in the Workplace

Mental Health America: Workplace Information

National Partnership for Workplace Mental Health

Health and Human Development Warning Signs in the Emergency Department

Strategies for Suicide Prevention in Veterans

Working Together To Serve Veterans