# TUBERCULOSIS IN NASSAU COUNTY 2023

Tuberculosis, also known as TB, is a reportable and contagious bacterial disease. Most commonly, it affects the lungs. However, it can also affect other parts of the body, such as the lymph nodes, bones, joints, and brain. It is characterized by chronic cough, fever, weight loss, and night sweats. Pulmonary TB is spread through the air when someone with the disease coughs, sneezes, speaks, or sings. TB can be treated through a combination of several different medications taken for at least 6 months. Nassau County had a total of 45 TB cases in 2023. In 2023, the rate of TB in Nassau County was 3.22 cases per 100,000 individuals, which increased from 2022's rate of 2.94 cases per 100,000 individuals.

## **DEMOGRAPHICS**

### AGE

- Most TB cases in Nassau County occurred in patients between the ages of 45-64, 17 cases (38%), a rate of 4.4 per 100,000 population.
- The highest rate of TB occurred in the 65+ population, 5.5 per 100,000 population.
- The average age for all patients was 55.



- Males accounted for 58% of all county cases while females accounted for 42% of all county cases.
- Males had a higher rate of TB than females with a rate of 3.8 cases per 100,000 compared to 2.6 cases per 100,000.




Figure 1: Percentage and rate of all tuberculosis cases by age group, Nassau County 2023

### BIRTHPLACE

 89% of patients with TB disease were foreign-born and 11% were U.S. born in 2023.




Figure 2: Tuberculosis cases by sex, Nassau County, 2023

### **CASES BY BIRTHPLACE**




Figure 3: Tuberculosis cases of foreign-born and U.S.-born, Nassau County, 2023





## CASES BY SEX

## **DISEASE DATA**

### SITE OF DISEASE

TB can be pulmonary (occurring in the lungs), extrapulmonary (occurring somewhere other than the lungs), or a combination of the two.


- The majority, 64.4%, of all cases of TB were pulmonary only.
- o 13.3% were both pulmonary and extrapulmonary.
- o 22.2% of cases were extrapulmonary only.

#### PULMONARY

Individuals suspected of having pulmonary TB have sputum samples collected for AFB smear and culture tests. A positive AFB smear is a strong indication that a patient is infectious. A patient that tests negative for AFB smear can still have a positive sputum culture, which confirms the diagnosis of TB disease. Culture conversion is used to determine treatment success.

- o 37.8% of all TB cases were sputum AFB smear positive.
- o 55.6% of all TB cases were sputum culture positive.
- 86.4% of sputum culture-positive cases converted to negative within 60 days of treatment initiation.

### SITE OF DISEASE



Pulmonary Only

- Pulmonary and Extrapulmonary
- Extrapulmonary Only

Figure 4: Tuberculosis cases by disease site, Nassau County, 2023




Figure 5: Extrapulmonary cases by disease site, Nassau County, 2023

### LATENT TUBERCULOSIS INFECTION

Latent TB Infection (LTBI) occurs when someone is infected with TB bacteria but does not have active TB. People with LTBI do not have any symptoms and are not infectious. They can, however, develop TB disease if the bacteria are activated. Generally, 5-10% of those with LTBI develop TB disease (CDC, 2014).

- Nassau County Department of Health identified 229 contacts to active TB cases. Of those, 58.5% were evaluated (134).
- 23.9% of evaluated contacts were diagnosed with LTBI (32).
- Of those diagnosed with LTBI, 78.1% started LTBI treatment (25), 32.0% of those completed LTBI treatment (8), 52.0% are still being treated (13), and 20.0% stopped LTBI treatment (5).

#### TREATMENT

Directly Observed Therapy (DOT) is the most effective way to ensure patients adhere to and complete their treatments. The goal is for patients to complete treatment within 12 months. DOT is offered by the Department of Health's TB Control Bureau to all patients receiving treatment for TB disease.

- o 75.6% of eligible patients received DOT administered by the Nassau County Department of Health or another facility.
- o Of those eligible to complete treatment, 80.0% completed treatment within 12 months.

